

Pro

LEBANON Scape®

**FERTILIZER
WITH TRIMEC®
PLUS DIMENSION®
WEED CONTROL**

18-0-6

MESA

P A T E N T E D

GUARANTEED ANALYSIS

Total Nitrogen (N) 18%
2.5% Ammoniacal Nitrogen
1.6% Water Insoluble Nitrogen*
11.3% Urea Nitrogen
2.6% Other Water Soluble Nitrogen*
Soluble Potash (K₂O) 6%
Sulfur (S) 2.8%

Derived From: Ammonium Sulfate, Methylene Ureas, Urea, Muriate of Potash.

*4.2% Slowly available Nitrogen from methylene urea.

Chlorine (Cl) not more than 5.0%

NOTE: Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 lb. Nitrogen per 1,000 sq. ft.

SUGGESTED SPREADER SETTINGS			
Spreader	3.6#/1,000 sq. ft.	Spreader	3.6#/1,000 sq. ft.
LebanonTurf	4½	Lesco (letter/Numeric)	i/17
Andersons AccuPro	L	PennMulch HVO	M
Earthway Rotary	15	ProScape SS	M
Gandy	27	Spyker	4½
Lely	5½ II	Vicon (all models)	25

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation, and evenness of terrain may require slightly different settings for desired coverage.

ProScape, LebanonTurf and PennMulch are registered trademarks of Lebanon Seaboard Corporation.

Information regarding the contents and levels of metals in this product is available on the Internet at: <http://www.regulatory-info-lebsea.com>

EPA Reg. No. 961-419

(L) (D)

EPA Est. No. 961-PA-01:961-IL-01

Superscript used is first letter of lot number.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for First Aid and additional Precautionary Statements.

MANTENGA ALEJADO DEL ALCANCE DE LOS NIÑOS

PRECAUCIÓN

Consulte el panel posterior para obtener información sobre primeros auxilios y otras declaraciones de precaución adicionales.

ACTIVE INGREDIENTS:

2,4-D, 2-ethylhexyl ester.....1.108%
Mecoprop-p 0.167%
Dicamba 0.071%
Dithiopyr 0.160%
OTHER INGREDIENTS:.....98.494%
TOTAL 100.000%

THIS PRODUCT CONTAINS:

14.70 lb 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.735% (CAS 1928-43-4)
3.34 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton or 0.167% (CAS 16484-77-8)
1.42 lb 3,6-dichloro-o-anisic acid equivalent per ton or 0.071% (CAS 1918-00-9)
3.20 lb dithiopyr per ton or 0.160% (CAS 97886-45-8)
Isomer Specific By AOAC Methods.

New York State: Do not exceed one application at the maximum rate of 3.6 pounds of product per 1,000 sq.ft. per year (0.25 lb. of dithiopyr per acre per year).



A/18

22-53894



Manufactured by:

Lebanon Seaboard Corporation

1600 E. Cumberland St. • Lebanon, PA 17042

Made in the U.S.A.

NET WEIGHT **40** LBS. (18.14 KG)

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

First Aid	
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 - 20 minutes.• Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of emergency, call toll free (888)-208-1368	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

• Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.

USE INSTRUCTIONS

This product is formulated to provide control of most annual grasses and certain broadleaf weeds* commonly found in home lawns. This product contains three components: a preemergent herbicide, a lawn fertilizer, and a post emergent broadleaf weed* killer.

• The preemergent herbicide can be applied to established turfgrass to control crabgrass goosegrass, foxtails, and other undesirable annual grasses.

• The fertilizer contains nutrients for turfgrass uptake.

• The post emergent broadleaf herbicide will control over 250 lawn weeds* including dandelion, chickweed, thistle and clover.

A list of broadleaf weeds is presented on this label.

*See use directions for complete list of treated weeds.

Controls 253 weeds!

Amaranth, purple	Charlie)	Mugwort	Speedwell, bi-lobed
Amaranth, slender	Cupid shaving Brush	Mustard, ball	Speedwell, common
Aster, heath	Cudweed	Mustard, black	Speedwell, corn
Aster, white heath	Daisy, English	Mustard, blue	Speedwell, ivyleaf
Aster, purple	Daisy, oxeye	Mustard, hedge	Speedwell, persian
Aster, white prairie	Dandelion, common	Mustard, Indian	Speedwell, purslane
Artichoke	Dandelion, false	Mustard, tansy	Speedwell, slender
Austrian fieldcress	Day flower	Mustard, tumble	Speedwell, snow
Bedstraw, Catchweed	Deadnettle	Mustard, white	Speedwell, thymeleaf
Bedstraw, smooth	Dichondra	Mustard, wormseed	Spiny amaranth
Bedstraw, Northern	Dock, broadleaf	Nettle, stinging (burning)	Spurge, spotted
Bedstraw, pineywoods	Dock, curly	Nettle, tall	Spurge, prostrate
Beggarticks, hairy	Dogbane	Nettle, wood	Spurweed
Betony, Florida	Dogfennel	Parsley-piert	Star of Bethlehem
Bindweed, Field	Dollarweed (Pennywort)	Parsnip	Strawberry, India mock
Bindweed, Hedge	Dove weed	Pearwort	Tansy ragwort
Bitter Wintercress	Elderberry	Pennycress (frenchweed)	Tanweed
Bittercress, hairy	Falseflax	Pepperweed, perennial	Thistle, blessed
Bitterweed	False sunflower	Pepperweed, Virginia	Thistle, bull
Black-eyed Susan	Fiddleneck	(Peppergrass)	Thistle, Canada
Black Medic	Filaree, redstem	Pigweed, prostate	Thistle, flodman (prairie)
Blood flower milkweed	Filaree, whitestem	Pigweed, redroot	Thistle, musk
Blue lettuce	Fleabane, daisy (annual)	Pigweed, Russian	Thistle, platte
Brassbuttons	Fleabane, rough	Pigweed, smooth	Thistle, plumeless
Bristly oxtongue	Galinsoga, hairy	Pigweed, tumble	Thistle, Russian
Broomweed	Galinsoga, smallflower	Pineapple weed	Thistle, Scotch
Buckhorn	Goldenrod	Plains coreopsis (tickseed)	Thistle, wavyleaf
Bullnettle	Ground ivy	Plantain, blackseed	Thistle, yellow (horrible)
Burclover	Groundsel	Plantain, bracted	Trailing crownvetch
Burdock	Gumweed	Plantain, broadleaf	Velvetleaf
Burweed, lawn (spurweed)	Hairy fleabane	Plantain, buckhorn	Vervain, blue
Buttercup, bulbous	Hawkweed, meadow	Plantain, hoary	Vervain, hoary
Buttercup, creeping	Hawkweed, mouseear	Plantain, narrowleaf	Vervain, prostrate
Buttonweed, common (poorjoe)	Hawkweed, orange	Plantain, slender	Vervain, tall
Buttonweed, Virginia	Hawkweed, yellow	Plantain, woolly	Vetch, bird
Carolina geranium (wild)	Healall	Poison ivy	Vetch, hairy
Carpetweed	Heartleaf drymary	Poison oak	Vetch, milk
Catnip	Hemp	Pokeweed	Vetch, narrowleaf
Catsear, spotted (false dandelion)	Henbit	Prairie Sunflower	Vetch, wild
Chamber bitter	Hoary cress	Prickly lettuce (compass plant)	Virginia creeper
Chickweed, common	Horsenettle	Prickly sida	Western clematis
Chickweed, mouseear	Horseweed	Prostrate knotweed	Western salsify
Chickweed, sticky	Jimsonweed	Puncturevine	Wild aster
Chicory	Johnny-Jump violet	Purslane, common	Wild buckwheat
Cinquefoil	Knawel	Pusley, Brazil	Wild carrot
Clover, crimson	Knotweed, prostrate	Pusley, Florida	Wild four-o'clock
Clover, alyce	Kochia	Pusley, large flower	Wild garlic
Clover, hop	Lambsquarters	Ragweed, bur	Wild lettuce
Clover, rabbit foot	Lespedeza	Ragweed, common	Wild marigold
Clover, red	Mallow, alkali	Ragweed, lanleaf	Wild mustard (yellow rocket)
Clover, strawberry	Mallow, bristly	Ragweed, western	Wild onion
Clover, yellow sweet	Mallow, common	Redstem filaree	Wild parsnip
Clover, white sweet	Mallow, dwarf	Rough cinquefoil	Wild radish
Clover, white	Mallow, venice	Scarlet pimpernel	Wild rape
Cockle	Marcela	Shepardspurse	Wild strawberry
Cocklebur	Matchweed	Smartweed, ladythumb	Wild sweet potato
Cocklebur, oriental	Mexicanweed	Smartweed, pale	Wild violet
Cocklebur, spiny	Moneywort	Smartweed, Pennsylvania	Woodsorrel, creeping
Common mullain	Morningglory, bigroot	Smooth chaff-flower	Woodsorrel, violet
Corn Spurry	Morningglory, ivyleaf	Smoothdock	Woodsorrel, yellow (Oxalis)
Creeping Beggarweed	Morningglory, mexican	Sorrel, red (sheep)	Woolly croton
Creeping Jenny (creeping	Morningglory, red	Sowthistle, annual	Wormseed
	Morningglory, tall	Sowthistle, spiny	Yarrow, common
	Morningglory, wolly	Spanish needles	Yarrow, western
			Yarrow, yellow

WHERE TO USE

Lebanon ProScape Fertilizer with Trimec Plus Dimension Weed Control may be used on the following established turf species:

Established Cool Season Turfgrass: Kentucky bluegrass, perennial ryegrass, fine fescues, tall fescue

Established Warm Season Turfgrass: bahiagrass, bermudagrass, buffalograss, zoysiagrass

Do not use this product on dichondra, carpetgrass or colonial bentgrass.

Do not use this product on St. Augustinegrass in Florida.

Do not use this product on turfgrass where desirable clovers are present.

If the lawn has been seeded recently, wait until after the third mowing before using this product. Do not seed, overseed or sprig the treated area for 10 weeks after application.

USE RESTRICTIONS

The use of this product on lawns and ornamental turfgrasses that are not well established, or have been weakened by stress due to mechanical operations, chemical injury, disease, insects, pests, excessive temperatures, or lack of moisture may increase the chances for turfgrass injury. Allow turfgrass weakened by these stresses to recover before application of this product.

Do not use this product on or near desirable plants, including within the drip line of the roots of desirable trees and shrubs, since injury may result.

Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches.

Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans.

Do not apply when windy.

Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the

treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

Do not exceed specified dosage for any area.

Do not use clippings from treated areas for composting or for mulching vegetables or fruit trees.

WHEN TO APPLY

This product may be applied to established lawns as a single application or as split applications in the spring, early summer, or fall.

For the best results in the spring, apply this product one to two (1 to 2) weeks before the expected germination of crabgrass. When the period of germination for crabgrass and other weeds listed on this label extends into early summer, then a second application can be made at 6 to 10 weeks after the initial application.

For control of crabgrass and other listed grasses: This product is a preemergent herbicide to control crabgrass (smooth, southern and large) and other susceptible annual grasses. This product may also provide early post emergent control of crabgrass. However, post emergent crabgrass control will be obtained only when this product is applied before the first tiller stage of crabgrass.

For broadleaf weed control: Best results are obtained during the spring and fall when broadleaf weeds are present, young and actively growing under adequate soil moisture and favorable temperatures (50°to 80°F).

HOW TO APPLY

This product needs to stick to the leaves of broadleaf weeds and should be applied when the grass is wet from dew or rain; poor results will occur if it is applied to dry lawns. The lawn should not be watered or disturbed for 24 hours after application. Delay mowing for 3 days after application.

Apply this product when the wind is calm. Avoid spreading granules onto flowers, vegetables, and ornamental shrubbery or desirable plants.

HOW MUCH TO APPLY

Kentucky bluegrass, fescue spp., perennial ryegrass, common bermudagrass, and zoysiagrass. Apply 3.6 pounds of product per 1,000 square feet of established turfgrass.

Hybrid bermudagrass, bahiagrass and buffalograss. Apply 3 pounds of product per 1,000 square feet of established turfgrass.

New York State: Do not exceed one application at the maximum rate of 3.6 pounds of product per 1,000 sq. ft. per year (0.25 lb. active ingredient of dithiopyr per acre per year).

Other States: The maximum broadcast application rate is 3.6 pounds of product per 1,000 sq. ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 7.2 pounds of product per 1,000 sq.ft.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Keep in original container in a cool, dry area, away from extreme heat and direct sunlight.

Do not store near food feed, seeds or other pesticides.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container.

If empty: Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.